

Applicant : Jacob Bar-Tana  
Serial No. : 10/735,439  
Filed : December 11, 2003  
Page 2 of 3: Communication in Response to November 28, 2008 Notice

**REMARKS**

In the November 28, 2008 Notice, a copy of which is attached hereto as **Exhibit A**, it was stated that the publications contained within "Exhibit G" and "Exhibit I" of the Amendment filed on August 28, 2008 (actually filed August 25, 2008), were unclear and illegible.

In response, applicant submits as **Exhibit B** and **Exhibit C** attached hereto replacement copies of the publications contained within "Exhibit G" and "Exhibit I" of the Amendment filed on August 25, 2008 respectively.

If a telephone interview would be of assistance in advancing prosecution of the subject application, the undersigned attorney invites the Examiner to telephone him at the telephone number provided below.

Applicant : Jacob Bar-Tana


Serial No. : 10/735,439

Filed : December 11, 2003

Page 3 of 3: Communication in Response to November 28, 2008 Notice

No fee, other than the enclosed \$245.00 fee for a two-month extension of time is deemed necessary in connection with the filing of this Communication. However, if any additional fee is required, authorization is hereby given to charge the amount of any such fee to Deposit Account No. 03-3125.


Respectfully submitted,




John P. White  
Registration No. 28,678  
Attorney for Applicant  
Cooper & Dunham LLP (Customer #23432)  
30 Rockefeller Plaza  
20<sup>th</sup> Floor  
New York, New York 10112  
(212) 278-0400

I hereby certify that this correspondence is being deposited on this date with the U.S. Postal Service with sufficient postage as first class mail in an envelope addressed to:

Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450.

 2/27/09  
John P. White Date  
Reg. No. 28,678

# **EXHIBIT A**



UNITED STATES DEPARTMENT OF COMMERCE

U.S. Patent and Trademark Office

Address: COMMISSIONER FOR PATENTS  
P.O. Box 1450  
Alexandria, Virginia 22313-1450

APPLICATION NO./ CONTROL NO.	FILING DATE	FIRST NAMED INVENTOR / PATENT IN REEXAMINATION	ATTORNEY DOCKET NO.
10/735,439	12/11/2003	BAR-TANA, JACOB	1567/70937-ZA /JPW/AG

John P. White  
Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, NY 10036

*Reply Due: 12/28/08*

EXAMINER

Leslie A. Royds

ART UNIT	PAPER
----------	-------

1614

20081121

DATE MAILED:

**Please find below and/or attached an Office communication concerning this application or proceeding.**

Commissioner for Patents

NOTICE OF NON-RESPONSIVE AMENDMENT

Applicant's amendment and remarks filed August 28, 2008 have each been received and entered into the present application.

Consideration of the remarks and documents contained therein has been performed herein by the Examiner. It is noted that Applicant discusses and relies upon the content designated as "Exhibit G" and "Exhibit I" to establish and support Applicant's position that the instant claims circumscribe patentable subject matter over the cited prior art. However, it is noted on the record that each of these publications upon which Applicant relies is sufficiently unclear to preclude a full and complete consideration of the information contained therein to support Applicant's position that the claimed subject matter distinguishes over the prior art cited under 35 USC 103(a).

Accordingly, by issuance of this notice, Applicant is requested (and afforded the opportunity) to provide a full, clear and legible copy of the publications contained within "Exhibit G" and "Exhibit I" as relied upon in the Remarks for consideration by the Examiner of record in order to fully and properly consider the persuasiveness of Applicant's remarks regarding the rejections of record.

Since the above-mentioned response appears to be a bona fide attempt to reply, Applicant is given a TIME PERIOD of ONE (1) MONTH or THIRTY (30) DAYS, whichever is longer, from the mailing date of this notice within which to supply the omission or correction, in order to avoid abandonment. Extensions of this time period under 37 C.F.R. 1.136(a) are available.

Any inquiry concerning this communication or earlier communications from the Examiner should be directed to Leslie A. Royds, whose telephone number is (571)-272-6096. The Examiner can normally be reached from Monday through Friday, 9:00 AM to 5:30 PM. If attempts to reach the Examiner are unsuccessful, the Examiner's supervisor, Ardin H. Marschel, can be reached on (571)-272-6088. The fax phone number for the organization where this application or proceeding is assigned is (571)-273-8300.

/Ardin Marschel/  
Supervisory Patent Examiner, Art Unit 1614

/Leslie A. Royds/  
Patent Examiner, Art Unit 1614



# UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office  
Address: COMMISSIONER FOR PATENTS  
P.O. Box 1450  
Alexandria, Virginia 22313-1450  
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
10/735,439	12/11/2003	Jacob Bar-Tana	1567/70937-ZA /JPW/AG	2054

John P. White  
Cooper & Dunham LLP  
1185 Avenue of the Americas  
New York, NY 10036

7590

11/28/2008

EXAMINER
----------

ROYDS, LESLIE A

ART UNIT	PAPER NUMBER
----------	--------------

1614

MAIL DATE	DELIVERY MODE
-----------	---------------

11/28/2008

PAPER

**Please find below and/or attached an Office communication concerning this application or proceeding.**

The time period for reply, if any, is set in the attached communication.

## **EXHIBIT B**

SEVENTEENTH EDITION

# THE MERCK MANUAL OF DIAGNOSIS AND THERAPY

הספריה הרפואית הלאומית  
ע"ש מנחם פרידמן  
ת.ד. 12272, ירושלים 91120

*Editors*  
MARK H. BEERS, M.D., and ROBERT BERKOW, M.D.

*Senior Assistant Editors*  
ROBERT M. BOGIN, M.D., and  
ANDREW J. FLETCHER, M.B., B.Chir.

*Editorial Board*  
Philip K. Bondy, M.D.  
Preston V. Dilts, Jr., M.D.  
Douglas A. Drossman, M.D.  
L. Jack Faling, M.D.  
Eugene P. Frenkel, M.D.  
Glen O. Gabbard, M.D.  
Robert A. Hoekelman, M.D.  
Gerald L. Mandell, M.D.  
Fred Plum, M.D.  
G. Victor Rossi, Ph.D.  
Paul H. Tanser, M.D., F.R.C.P.(C)

*Published by* MERCK RESEARCH LABORATORIES

*Division of* MERCK & Co., Inc.  
Whitehouse Station, N.J.

1999

EDITIONS

vices, Cyprus  
ishing House, Beijing

GmbH, Munich  
lishing, Athens

an  
law  
São Paulo  
ow

CS

MANUAL

GERIATRICS

MEDICAL INFORMATION—HOME EDITION

a nonprofit basis as a service to the scientific com-

### **Editorial and Production Staff**

*Executive Editor:* Keryn A.G. Lane

*Senior Staff Editor:* Susan T. Schindler

*Staff Editor:* Julie Kostecky

*Staff Editor:* Sandra J. Masse

*Production Editor:* Debra G. Share

*Contributing Editor:* Roger I. Schreck, M.D.

*Designer:* Lorraine B. Kilmer

*Indexer:* Susan Thomas, Ph.D.

*Textbook Production Coordinator:* Diane C. Zenker

*Medical Textbook Coordinator:* Dorothy A. Bailey

*Executive Assistant:* Diane Cosner-Bobrin

*Publisher:* Gary Zelko

*Advertising and Promotional Supervisor:* Pamela J. Barnes

Library of Congress Catalog Card Number 1-31760

ISBN 0911910-10-7

ISSN 0076-6526

Copyright 1999 by Merck & Co., Inc.

All rights reserved. No part of this book may be reproduced or used in any form or by any means, electronic or mechanical, including photocopying, or by any information storage and retrieval system, without permission in writing from the Publisher. Inquiries should be addressed to The Merck Manuals Department, P.O. Box 4, Merck & Co., West Point, PA 19486.

Printed in England by Clays Ltd, St Ives plc



TABLE 296-2. NORMAL LABORATORY VALUES (Continued)

Test	Specimen	Method	Normal Range	
			CONVENTIONAL UNITS	SI UNITS
Bilirubin Direct Indirect	Serum Serum	Colorimetry Colorimetry Calculation: Total bilirubin minus direct bilirubin	$\leq 0.4$ mg/dL $\leq 1.3$ mg/dL	$\leq 7$ $\mu$ mol/L $\leq 22$ $\mu$ mol/L
Total	Serum	Colorimetry	$\leq 1.3$ mg/dL	$\leq 22$ $\mu$ mol/L
Bleeding time, template	Not appli- cable	Template method	2.6–9.5 min	
CA 15-3	Serum	ABBOTT AXSYM CA 15-3 MEIA	$< 32$ U/mL	$< 32$ kU/L
CA 19-9*	Serum	CIS ELISA-CA 19-9 IRMA	$< 33$ U/mL	$< 33$ kU/L
CA 27.29	Serum	BIOMIRA TRUQUANT BR RIA	$< 38$ U/mL	$< 38$ kU/L
CA 125	Serum	CENTOCOR CA 125 II RIA	$< 35$ U/mL	$< 35$ kU/L
Cadmium	Blood	Atomic spectroscopy	$\leq 2$ $\mu$ g/L (nonsmoker)	$\leq 17.8$ nmol/L (nonsmoker)
Calcitonin	Serum	Imunoassay	Male: $< 13.8$ pg/mL Female: $< 6.4$ pg/mL	Male: $< 13.8$ ng/L Female: $< 6.4$ ng/L
Calcium	Serum Urine	Colorimetry Colorimetry	8.5–10.3 mg/dL Male: $< 300$ mg/day Female: $< 250$ mg/day	2.12–2.57 mmol/L Male: $< 7.5$ mmol/day Female: $< 6.2$ mmol/day
Carbon dioxide	Serum	Colorimetry	20–32 mmol/L	20–32 mmol/L
Carboxyhemoglobin	Blood	Spectrophotometry	$< 2\%$ of total Hb (nonsmoker)	$< 0.02$
Carcinoembryonic antigen (CEA)	Serum	CHIRON ACS:180 ICMA	$< 2.5$ ng/mL (nonsmoker)	$< 2.5$ $\mu$ g/L (nonsmoker)
Carotene	Serum	Colorimetry	60–300 $\mu$ g/dL	0.9–5.6 $\mu$ mol/L
<b>Catecholamines</b>				
		Plasma	HPLC	
		Dopamine	Supine: $< 90$ pg/mL Standing: $< 90$ pg/mL	Dopamine Supine: $< 588$ pmol/L Standing: $< 588$ pmol/L
		Epinephrine	Supine: $< 50$ pg/mL Standing: $< 90$ pg/mL	Epinephrine Supine: $< 273$ pmol/L Standing: $< 491$ pmol/L
		Norepinephrine	Supine: 110–410 pg/mL Standing: 125–700 pg/mL	Norepinephrine Supine: 650–2423 pmol/L Standing: 739–4137 pmol/L
		Plasma	HPLC	
		Supine	120–450 pg/mL Standing: 150–750 pg/mL	Supine: 709–2860 pmol/L Standing: 887–4433 pmol/L
Total	Plasma	HPLC	25–63 ng/dL	250–630 ng/L
Ceruloplasmin	Serum	Nephelometry	95–108 mg/dL	95–108 nmol/L
Chloride	Serum	ISE	Desirable: $< 200$ mg/dL Borderline-high: 200–239 mg/dL High: $\geq 240$ mg/dL	Desirable: $< 5.17$ mmol/L Borderline-high: 5.17–6.18 mmol/L High: $\geq 6.21$ mmol/L
Cholesterol, total	Serum	Colorimetry	75–161 mg/dL 16–47 mg/dL 31–66 U/mL	0.75–1.61 g/L 0.16–0.47 g/L 31–66 kU/L
<b>Complement</b>				
C3	Serum	Nephelometry		
C4	Serum	Nephelometry		
Total (CH <sub>50</sub> )	Serum	Liposome lysis		
Complete blood count (CBC)	Blood	Automated hematology analyzer		

CA 15-3	Serum	ABBOTT AXSYM CA 15-3 MEIA	< 32 U/mL
CA 19-9*	Serum	CIS ELISA-CA 19-9 IRMA	< 33 U/mL
CA 27.29	Serum	BIOMIRA TRUQUANT BR RIA	< 38 U/mL
CA 125	Serum	CENTOCOR CA 125 II RIA	< 35 U/mL
Cadmium	Blood	Atomic spectroscopy	≤ 2 µg/L (nonsmoker)
Calcitonin	Serum	Immunoassay	Male: < 13.8 pg/mL Female: < 6.4 pg/mL
Calcium	Serum Urine	Colorimetry Colorimetry	8.5-10.3 mg/dL Male: < 300 mg/day Female: < 250 mg/day
Carbon dioxide	Serum	Colorimetry	20-32 mmol/L
Carboxyhemoglobin	Blood	Spectrophotometry	< 2% of total Hb (nonsmoker)
Carcinoembryonic antigen (CEA)	Serum	CHIRON ACS:180 ICMA	< 2.5 ng/mL (nonsmoker)
Carotene	Serum	Colorimetry	50-300 µg/dL

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Abbott Diagnostics

Dopamine	Supine: < 90 pg/mL Standing: < 90 pg/mL	Dopamine	Supine: < 588 pmol/L Standing: < 588 pmol/L
Epinephrine	Supine: < 60 pg/mL Standing: < 90 pg/mL	Epinephrine	Supine: < 273 pmol/L Standing: < 491 pmol/L
Norepinephrine	Supine: 110-410 pg/mL Standing: 125-700 pg/mL	Norepinephrine	Supine: 650-2423 pmol/L Standing: 739-4137 pmol/L
Supine: 120-450 pg/mL Standing: 150-750 pg/mL		Supine: 709-2650 pmol/L Standing: 887-4433 pmol/L	
25-63 mg/dL		250-630 mg/L	
95-108 mmol/L		96-108 mmol/L	
Desirable: < 200 mg/dL Borderline-high: 200-238 mg/dL High: ≥ 240 mg/dL		Desirable: < 5.17 mmol/L Borderline-high: 5.17-6.18 mmol/L High: ≥ 6.21 mmol/L	
75-161 mg/dL 16-47 mg/dL 31-66 U/mL		0.75-1.61 g/L 0.16-0.47 g/L 31-66 kU/L	
Male: 13.8-17.2 g/dL Female: 12.0-15.6 g/dL Male: 41-50% Female: 35-46% Male: 4.4-5.8 × 10 <sup>9</sup> /µL Female: 3.9-5.2 × 10 <sup>9</sup> /µL Mean corpuscular volume: 78-102 fL Mean corpuscular Hb: 27-33 pg Mean corpuscular Hb concentration: 32-36 g/dL RBC distribution width: ≤ 15% 3.8-10.8 × 10 <sup>9</sup> /µL Absolute neutrophils: 1500-7800 cells/µL Absolute eosinophils: 50-550 cells/µL		Male: 138-172 g/L Female: 120-158 g/L Male: 0.41-0.50 Female: 0.35-0.46 Male: 4.4-5.8 × 10 <sup>9</sup> /L Female: 3.9-5.2 × 10 <sup>9</sup> /L Mean corpuscular volume: 78-102 fL Mean corpuscular Hb: 27-33 pg Mean corpuscular Hb concentration: 32-36 g/L RBC distribution width: ≤ 0.15 3.8-10.8 × 10 <sup>9</sup> /L Absolute neutrophils: 1.5-7.8 × 10 <sup>9</sup> /L Absolute eosinophils: 0.05-0.55 × 10 <sup>9</sup> /L	

Table continues on the following page.

Test	Specimen	Method	Normal Range	81 Units
Human chorionic gonadotropin (hCG)				
Qualitative	Urine	Immunoassay	Nonpregnant: negative	
Quantitative (intact and free $\beta$ )	Serum	Immunoassay	Pregnant: positive Male: < 2 IU/L Female: Prenenopausal: < 5 IU/L Postmenopausal: < 10 IU/L Pregnancy: 0-2 wk < 500 IU/L 2-3 wk 100-5,000 IU/L 3-4 wk 500-10,000 IU/L 1-2 mo 1,000-200,000 IU/L 2-3 mo 10,000-100,000 IU/L Male: 8.3-41.4 $\mu$ mol/day Female: 6.6-33.1 $\mu$ mol/day 3-47 $\mu$ mol/day	
17-Hydroxycorticosteroids	Urine	Enzymatic colorimetry	Male: 3-15 mg/day Female: 2-12 mg/day	
5-Hydroxyindoleacetic acid (5-HIAA)	Urine	HPLC	0.5-9.0 mg/day	
Immunoglobulin				
IgA	Serum	Nephelometry	81-463 mg/dL $\leq 14$ mg/dL	
IgD	Serum	Radial immunodiffusion	< 180 U/mL	
IgE	Serum	Immunoassay	Subclass IgG 1: 450-900 mg/dL Subclass IgG 2: 180-530 mg/dL Subclass IgG 3: 13-80 mg/dL Subclass IgG 4: 8-100 mg/dL	
IgG, subclasses	Serum	Nephelometry	723-1686 mg/dL 48-271 mg/dL	
IgG, total	Serum	Nephelometry	5-25 $\mu$ U/mL	
IgM	Serum	Nephelometry	26-170 $\mu$ g/dL	
Inulin	Serum	Immunoassay		
Iron	Serum	Colorimetry		
Iron-binding capacity	Serum	Colorimetry	200-450 $\mu$ g/dL % Saturation: 12-57%	
17-Ketogenic steroids	Urine	Colorimetry	Male: 5-23 mg/day Female: 3-15 mg/day	
17-Ketosteroids, total	Urine	Colorimetry	Male: 9-22 mg/day Female: 5-15 mg/day	
Lactate dehydrogenase (LD)				
Isoenzymes	Serum	Electrophoresis	LD1: 20-36% of total LD2: 32-50% of total LD3: 15-25% of total LD4: 2-10% of total LD5: 3-13% of total $\leq 270$ U/L	
Total	Serum	Enzymatic colorimetry	9-16 mg/dL	
Lactic acid	Plasma (venous)	Enzymatic colorimetry	< 25 $\mu$ g/dL	
Lead	Blood	Atomic spectroscopy	7-60 U/L	
Uryase	Serum	Enzymatic		

17-Hydroxycorticosteroids	Urine	Enzymatic colorimetry	2-3 wk 100-5,000 IU/L 3-4 wk 500-10,000 IU/L 1-2 mo 1,000-200,000 IU/L 2-3 mo 10,000-100,000 IU/L Male: 8.3-41.4 µmol/day Female: 5.6-33.1 µmol/day 3-47 µmol/day
5-Hydroxyindoleacetic acid (5-HIAA)	Urine	HPLC	
Immunoglobulin	Serum	Nephelometry	81-459 mg/dL ≤ 14 mg/dL
IgA	Serum	Radial immunodiffusion	< 180 U/mL
IgE	Serum	Immunosay	Subclass IgG 1: 450-900 mg/dL Subclass IgG 2: 180-520 mg/dL Subclass IgG 3: 13-80 mg/dL Subclass IgG 4: 8-100 mg/dL
IgG, subclasses	Serum	Nephelometry	723-1685 mg/dL
IgG, total	Serum	Nephelometry	48-271 mg/dL
IgM	Serum	Nephelometry	5-25 µU/mL
Inulin	Serum	Immunosay	25-170 µg/dL
Iron	Serum	Colorimetry	36-81 µmol/L % Saturation: 0.12-0.57

Iron-binding capacity	Serum	Colorimetry	200-450 µg/dL % Saturation: 12-57%
17-Ketogenic steroids	Urine	Colorimetry	Male: 6-23 mg/day Female: 3-15 mg/day
17-Ketosteroids, total	Urine	Colorimetry	Male: 9-22 mg/day Female: 5-15 mg/day
Lactate dehydrogenase (LD) isoenzymes	Serum	Electrophoresis	LD1: 20-36% of total LD2: 32-60% of total LD3: 16-25% of total LD4: 2-10% of total LD5: 3-13% of total ≤ 270 U/L
Total	Serum	Enzymatic colorimetry	9-16 mg/dL
Lactic acid	Plasma (venous)	Enzymatic colorimetry	< 25 µg/dL
Lead	Blood	Atomic spectroscopy	7-60 U/L
Lipase	Serum	Enzymatic colorimetry	Desirable: < 130 mg/dL Borderline-high: 130-169 mg/dL High: ≥ 160 mg/dL
Low density lipoprotein (LDL) cholesterol, direct	Serum	Immunoseparation/colorimetry	Male: 20-70 yr: 1.3-12.9 IU/L > 70 yr: 11.3-56.4 IU/L
Luteinizing hormone (LH)	Serum	Immunosay	Female: Follicular phase 0.8-25.8 IU/L Midcycle 25.0-57.3 IU/L Luteal phase 0.8-27.1 IU/L Pregnancy < 1.4 IU/L Postmenopausal 5.0-52.3 IU/L

Table continues on the following page.

## **EXHIBIT C**

## Brief Reports

## Nicotinic Acid as Therapy for Dyslipidemia in Non-Insulin-Dependent Diabetes Mellitus

Abhimanyu Garg, MBBS, MD, Scott M. Grundy, MD, PhD

Recently, nicotinic acid has been recommended as a first-line hypolipidemic drug. To determine the effectiveness of nicotinic acid in dyslipidemic patients with non-insulin-dependent diabetes mellitus, 13 patients were treated in a randomized crossover trial. Patients received either nicotinic acid (1.5 g three times daily) or no therapy (control period) for 8 weeks each. Compared with the control period, nicotinic acid therapy reduced the plasma total cholesterol level by 24%, plasma triglyceride level by 45%, very-low-density lipoprotein cholesterol level by 58%, and low-density lipoprotein cholesterol level by 15%, and it increased the high-density lipoprotein cholesterol level by 34%. However, nicotinic acid therapy resulted in the deterioration of glycemic control, as evidenced by a 16% increase in mean plasma glucose concentrations, a 21% increase in glycosylated hemoglobin levels, and the induction of marked glycosuria in some patients. Furthermore, a consistent increase in plasma uric acid levels was observed. Therefore, despite improvement in lipid and lipoprotein concentrations, because of worsening hyperglycemia and the development of hyperuricemia, nicotinic acid must be used with caution in patients with non-insulin-dependent diabetes mellitus with dyslipidemia. We suggest that the drug not be used as a first-line hypolipidemic drug in patients with non-insulin-dependent diabetes mellitus.

(JAMA. 1990;264:723-726)

**DYSLIPIDEMIA** is a common finding in non-insulin-dependent diabetes mellitus (NIDDM)<sup>1</sup> and probably contributes causally to coronary heart disease, a major cause of death in patients with NIDDM.<sup>2</sup> Recently, the National Cholesterol Education Program proposed new guidelines for the management of high blood cholesterol levels.<sup>3</sup> The guidelines obviously could not consider in depth every subgroup of patients with hyperlipidemias, and, therefore, problems of management of lipid and lipoprotein abnormalities in patients with NIDDM were not addressed in detail. The National Cholesterol Education Program recommended nicotinic acid and bile acid binding resins as first-line drugs for treatment of hypercholesterolemia, and nicotinic acid was designated as the drug of choice for hypercholesterolemic patients with concurrent hypertriglyceridemia. Since hypertriglyceridemia is the most prevalent lipid abnormality in NIDDM, the guidelines could be interpreted to mean that nicotinic acid is the drug of choice for diabetic dyslipidemia. Patients with NIDDM, however, have other metabolic abnormalities, and, therefore, the choice of hypolipidemic drug may not be the same as in nondiabetic patients. The purpose of this study was to examine the potential usefulness of nicotinic acid for the treatment of dyslipidemia in patients with NIDDM.

percholesterolemic patients with concurrent hypertriglyceridemia. Since hypertriglyceridemia is the most prevalent lipid abnormality in NIDDM, the guidelines could be interpreted to mean that nicotinic acid is the drug of choice for diabetic dyslipidemia. Patients with NIDDM, however, have other metabolic abnormalities, and, therefore, the choice of hypolipidemic drug may not be the same as in nondiabetic patients. The purpose of this study was to examine the potential usefulness of nicotinic acid for the treatment of dyslipidemia in patients with NIDDM.

## PATIENTS AND METHODS

## Patients

Thirteen male patients with NIDDM from a lipid clinic and a diabetes clinic were studied at the Veterans Administration Medical Center, Dallas, Tex. All patients had an insidious onset of diabetes after age 38 years, and none had a history of ketosis. Their ages ranged from 49 to 68 years (mean  $\pm$  SEM,  $59 \pm 1$  years). Body weights and body-mass indexes averaged  $91.7 \pm 3.3$  kg and

$29.9 \pm 0.7$  kg/m<sup>2</sup>, respectively. Four patients were receiving glyburide therapy, eight patients were receiving a combination of isophane insulin suspension and regular human insulin (Squibb-Novo, Princeton, NJ) subcutaneously before breakfast and supper for glycemic control, and one patient was receiving dietary therapy only. C-peptide levels were determined for patients receiving insulin therapy, both in the fasting state and 90 minutes after they ingested 480 mL of Sustacal (Mead Johnson & Co, Evansville, Ind); average values were  $751 \pm 182$  and  $1388 \pm 192$  pmol/L, respectively, confirming the diagnosis of NIDDM.<sup>4</sup> At entry, all patients had a plasma cholesterol level of 5.2 mmol/L or greater and/or a plasma triglyceride level of 2.8 mmol/L or greater. Six patients had coronary heart disease but none had recent myocardial infarction, unstable angina pectoris, or congestive heart failure. Patients were excluded if they had a history of peptic ulcer or gout, evidence of hyperuricemia (plasma uric acid concentration  $>475$   $\mu$ mol/L), or abnormal test results for liver, kidney, or thyroid gland functions. For patients taking specific hypolipidemic drugs, such therapy was discontinued at least 2 months prior to the study.

## Experimental Design

The study protocol was approved by the institutional review board, and each patient gave informed consent. All patients were studied during three hospitalizations in the metabolic ward, each lasting 5 days. Before being randomized, patients were hospitalized for 5 days, called the baseline period, during which the dosage of insulin or glyburide was adjusted to achieve good glycemic control, and energy intake was determined to project a constant body weight. Thereafter, no changes in the dosage of insulin or glyburide were allowed except to prevent symptomatic hypoglycemia. The plasma glucose con-

From the Veterans Administration Medical Center, Dallas (Drs Garg and Grundy), and the Center for Human Nutrition (Drs Garg and Grundy) and the Department of Clinical Nutrition (Drs Garg and Grundy), Internal Medicine (Drs Garg and Grundy), and Biochemistry (Dr Grundy), University of Texas Southwestern Medical Center at Dallas.

Reprint requests to Center for Human Nutrition, 5323 Harry Hines Blvd, Dallas, TX 75235-9052 (Dr Grundy).

JAMA, August 8, 1990—Vol 264, No. 6

Nicotinic Acid for Dyslipidemia in Diabetes—Garg &amp; Grundy 723




Fig 1.—Plasma levels of total cholesterol, low-density lipoprotein (LDL) cholesterol, triglycerides, very-low-density lipoprotein (VLDL) cholesterol, and high-density lipoprotein (HDL) cholesterol and the ratio of total to HDL cholesterol during the control and the nicotine acid periods in 13 patients with non-insulin-dependent diabetes mellitus with dyslipidemia. Each circle represents the mean of five daily determinations. Solid circles indicate mean values in patients receiving insulin therapy; open circles, values in patients receiving glyburide therapy or diet alone.

concentration was measured at 3, 7, and 11 AM and 4 and 9 PM each day. Fasting blood samples were drawn daily for analysis of lipids and lipoproteins. Blood was also drawn for a glycosylated hemoglobin determination and a routine hematologic and chemistry profile, including the uric acid concentration. Patients were instructed to follow an isocaloric diet throughout the study, the diet containing 50% carbohydrates, 30% fat, and 20% protein, with 300 mg of cholesterol.\* They were instructed not to consume alcohol during the trial.

After the baseline hospitalization, patients were randomized to receive nicotine acid or no therapy for a period of 8 weeks. All patients then crossed over to the drug/no therapy (control) period for the next 8 weeks. A double-blind, placebo-controlled trial was not planned because of previous reports of the ineffective nature of this design due to symptomatic side effects of nicotine acid therapy.<sup>6</sup> The nicotine acid dosage

Table 1.—Effect of Nicotinic Acid Therapy on Plasma Lipid and Lipoprotein Levels\*

	Study Period			P†
	Baseline	Control	Nicotinic Acid	
Plasma cholesterol, mmol/L	6.71 ± 0.33	6.35 ± 0.28	4.82 ± 0.29	.0001
Plasma triglycerides, mmol/L	5.08 ± 0.64	4.46 ± 0.62	2.45 ± 0.40	.0006
VLDL cholesterol, mmol/L	2.57 ± 0.40	2.19 ± 0.38	0.91 ± 0.19	.0006
LDL cholesterol, mmol/L	3.39 ± 0.42	3.40 ± 0.35	2.89 ± 0.21	.07
HDL cholesterol, mmol/L	0.76 ± 0.05	0.76 ± 0.04	1.02 ± 0.06	.0001
Total cholesterol/HDL cholesterol	9.22 ± 0.61	8.58 ± 0.55	4.93 ± 0.35	.0001

\*VLDL indicates very-low-density lipoprotein; LDL, low-density lipoprotein; and HDL, high-density lipoprotein. Values are mean ± SEM. To convert values from millimoles per liter to milligrams per deciliter, multiply the cholesterol values by 38.67 and the triglyceride values by 88.574.

†Comparison between the control and nicotine acid periods by a two-tailed paired *t* test.

was gradually increased from 50 mg three times daily on the first day to 1.5 g three times per day by the end of third week. Thereafter, patients continued to take the full dosage for the next 5 weeks. Patients reported as outpatients at two weekly intervals for a chemistry profile. On day 51 of each period, the patients entered the metabolic ward for 6 days, and blood samples were obtained each day as described above. On the last day of each period, plasma specimens were obtained every 2 hours for the determination of glucose levels. The patients were also interviewed about the side effects of the medication, such as flushing, rash, gastrointestinal distress, allergic reactions, and gout.

#### Biochemical Analyses

Fasting plasma samples were analyzed for total cholesterol, triglyceride, and lipoprotein cholesterol concentrations according to Lipid Research Clinic procedures,<sup>7</sup> except that cholesterol and triglyceride concentrations were measured enzymatically.<sup>8,9</sup> Briefly, very-low-density lipoprotein (VLDL, density <1.006 kg/L) was removed by preparative ultracentrifugation, and the cholesterol level was measured in the VLDL subfraction and the infranatant. The high-density lipoprotein (HDL) cholesterol level was measured in the supernatant after lipoproteins containing apolipoprotein B were precipitated by heparin-manganese.<sup>7</sup> Cholesterol in the low-density lipoprotein (LDL) fraction was taken as the difference between the cholesterol content of the 1.006-kg/L infranatant and HDL cholesterol.

The plasma glucose concentration was determined by glucose oxidase method using a glucose analyzer (Beckman Instruments Inc, Fullerton, Calif). Quantitative analysis of glycosylated hemoglobin was done by agar gel electrophoresis using kits (Helena Laboratories, Beaumont, Tex). The plasma C-peptide concentration was measured by

radioimmunoassay kits (Mallinckrodt Inc, St Louis, Mo).

#### Statistical Analyses

A repeated-measures analysis of variance test was performed to compare the baseline, nicotine acid, and control periods, to assess the effect of the sequence in which the patients were assigned to the control or active drug period, and to assess differences in response between patients receiving insulin and other therapy.<sup>10,11</sup> Multiple comparisons were made with use of the two-tailed paired *t* test with Bonferroni's correction. When three periods were included in the analysis,  $P < .0167$  was considered significant. The Wilcoxon signed rank test was used for data not consistent with the hypothesis of normality. The areas under the curve were compared with use of a *t* test. All results are expressed as mean ± SEM.

#### RESULTS

The analysis of variance did not reveal any differences in the response to nicotine acid therapy whether patients received insulin, glyburide, or no hypoglycemic drugs; therefore, plasma lipid and lipoprotein values in all patients were pooled. Results for each patient are shown in Fig 1, and results for all patients are summarized in Table 1. The order in which patients were allocated to the drug and control periods had no effect on the results. Plasma lipid and lipoprotein concentrations were not significantly different in the baseline and control periods (Table 1).

Compared with the control period, nicotine acid therapy reduced the plasma total cholesterol level by 24%. Plasma triglyceride levels were reduced by 45% ( $P < .001$ ) and VLDL cholesterol levels by 58% ( $P < .001$ ). The LDL cholesterol levels showed a modest 15% decrease with nicotine acid therapy which approached statistical significance ( $P = .07$ ). The HDL cholesterol concentrations rose consistently, with

Table 2.—Me

Mean plasma  
24-h plasma g  
mmol-HVLS  
insulin dosage  
glyburide dos  
glycosylated f  
24-h urinary gl  
Plasma uric ac  
Body weight, k

\*Values are  
†Compariso  
‡Plasma gl  
convert value  
§Plasma gl  
nine patients.  
||To convert |

an average  
The ratio  
cholestero  
during nic

The dail  
mic drugs  
patients.  
hypoglyce  
period, th  
duced by  
buride the  
persistent  
ported on  
control pe  
otinic ac  
pated max  
plasma gl  
of 8.2 mm  
to 18.2 an  
patient fo  
therapy),  
the daily i  
U. Despit  
insulin doe  
ma glucos  
tion remai  
during nic  
with value  
(7.8 mmol  
insulin do  
follow the  
excluded f

Overall,  
orated dur  
evidenced  
plasma gl  
9.1 mmol/l  
lated hemo  
ing nicotin  
glycosuria  
(Table 2 ar  
plasma glu  
of each pe  
revealed sig  
ing nicotin  
Nicotini  
ma uric a

Table 2.—Metabolic Variables During the Study\*

	Pt	Study Period			P†
		Baseline	Control	Nicotinic Acid	
Mean plasma glucose, mmol/L‡		7.19 ± 0.31	7.85 ± 0.50	9.09 ± 0.63	.068
24-h plasma glucose profile, mmol-H/L§			194.1 ± 6.2	213.5 ± 7.4	.047
Insulin dosage (n = 8), U/d		85.4 ± 9.5	87.4 ± 10.0	91.5 ± 10.2	.50
Glyburide dosage (n = 4), mg/d		6.25 ± 3.0	2.50 ± 1.0	6.25 ± 3.0	1.0
Glycosylated hemoglobin, %		9.8 ± 0.4	8.7 ± 0.6	10.5 ± 0.5	.002
24-h urinary glucose, g/d		3.1 ± 1.3	3.6 ± 1.4	14.9 ± 6.3	.016
Plasma uric acid, μmol/L		398 ± 22	382 ± 22	511 ± 33	.0001
Body weight, kg		91.7 ± 3.3	92.1 ± 2.8	91.1 ± 2.8	.19

\*Values are mean ± SEM.

†Comparison between the control and nicotinic acid periods by a two-tailed paired t test.

‡Plasma glucose values were determined at 3, 7, and 11 AM and 4 and 9 PM for 5 days during hospitalization. To convert values from millimoles per liter to milligrams per deciliter, multiply by 18.02.

§Plasma glucose values were determined at 2-hour intervals for 24 hours on the last day of hospitalization in nine patients. Values are given in area-under-the-curve units.

||To convert plasma uric acid values from micromoles per liter to milligrams per deciliter, multiply by 0.0168.

(Mallinckrodt

s analysis of need to compare id, and control effect of the se- tients were as- ctive drug per- ices in response ing insulin and le comparisons the two-tailed erroni's correc- s were included was considered on signed rank not consistent normality. The were compared results are ex-

nce did not re- he response to hether patients ide, or no hypo- re, plasma lipid in all patients or each patient l results for all in Table 1. The were allocated periods had no lasma lipid and ns were not sig- e baseline and

control period, duced the plas- l by 24%. Plas- ere reduced by DL cholesterol. The LDL cho- modest 15% de- acid therapy, tistical signifi- DL cholesterol sistent, with

an average increase of 34% ( $P < .0001$ ). The ratio of total cholesterol to HDL cholesterol also improved strikingly during nicotinic acid therapy.

The daily requirements of hypoglycemic drugs did not change in 10 of 13 patients. In one patient, due to mild hypoglycemic episodes in the control period, the daily insulin dosage was reduced by 4 U. In another patient, glyburide therapy was discontinued due to persistently low blood glucose levels reported on self-monitoring during the control period. In the third patient, nicotinic acid therapy caused an unanticipated marked deterioration in fasting plasma glucose values (from an average of 8.2 mmol/L during the control period to 18.2 and 21.5 mmol/L during the outpatient follow-up with nicotinic acid therapy), necessitating an increase in the daily insulin dosage from 76 to 105 U. Despite a 38% increase in the daily insulin dosage, the patient's mean plasma glucose values during hospitalization remained elevated (11.3 mmol/L) during nicotinic acid therapy compared with values during the control period (7.8 mmol/L). Although the increase in insulin dosage in this patient did not follow the original protocol, he was not excluded from analysis.

Overall, glycemic control deteriorated during nicotinic acid therapy, as evidenced by a 16% increase in mean plasma glucose levels, from 7.8 to 9.1 mmol/L. Concentrations of glycosylated hemoglobin increased by 21% during nicotinic acid therapy, and marked glycosuria was noted in some patients (Table 2 and Fig 2). A daylong profile of plasma glucose, obtained on the last day of each period in nine patients, also revealed significantly higher values during nicotinic acid therapy (Table 2).

Nicotinic acid therapy increased plasma uric acid levels in all the patients

(Table 2 and Fig 2). No patient, however, suffered from acute gouty arthritis. In two patients, mean plasma uric acid values rose to extremely high levels—684 and 761 μmol/L—with nicotinic acid therapy (Fig 2). Both of these patients had borderline low values of creatinine clearance, 1.05 and 1.08 mL/s, respectively, at entry into the study. A slight increase in the plasma creatinine concentration and a reduction in creatinine clearance was also noted in both patients during nicotinic acid therapy. No changes in the plasma creatinine concentration or creatinine clearance were noted in any other patients.

Most patients tolerated nicotinic acid therapy well except for minor complaints of flushing. None developed significant abnormalities in hepatic function test results throughout the study. One patient reported headaches but also noted improvement in claudication distance during nicotinic acid therapy. No patient dropped out as a consequence of side effects.

#### COMMENT

Soon after the discovery of the plasma lipid-lowering potential of nicotinic acid therapy by Altschul et al,<sup>12</sup> deterioration of glucose tolerance with this agent was reported in both nondiabetic subjects<sup>13,14</sup> and patients with NIDDM.<sup>15,17</sup> Since most of these claims were anecdotal, the potential clinical significance of this side effect has not been given due consideration. For instance, recent guidelines of the National Cholesterol Education Program can be taken to indicate that nicotinic acid is the drug of choice for treatment of dyslipidemia associated with NIDDM.<sup>18</sup> The current investigation, therefore, was carried out to examine carefully whether nicotinic acid will favorably modify plasma lipid and lipoprotein concentrations in pa-




Fig 2.—Mean plasma glucose, 24-hour urinary glucose, glycosylated hemoglobin, and plasma uric acid levels during the control and the nicotinic acid periods in 13 patients with non-insulin-dependent diabetes mellitus with dyslipidemia. Solid circles indicate mean values in patients receiving insulin therapy; open circles, values in patients receiving glyburide therapy or diet alone.

tients with NIDDM without significantly worsening their glycemic control.

In our patients, nicotinic acid therapy was highly effective for lowering levels of plasma triglycerides and VLDL cholesterol. It also raised levels of HDL cholesterol, with an increase averaging 34%. Total cholesterol levels were reduced significantly, as were total/HDL cholesterol ratios. Nicotinic acid therapy reduced LDL cholesterol levels in most but not all patients. Still, it generally *did not* produce the rise in LDL cholesterol concentrations commonly observed with other triglyceride-lowering therapies, eg, fibric acids<sup>19,20</sup> or n-3 polyunsaturated fatty acids.<sup>20,21</sup>

This study leaves little doubt that nicotinic acid therapy improves the lipoprotein profile in patients with NIDDM. On the other hand, the drug also causes a deterioration in glycemic control. In almost all patients, levels of glycosylated hemoglobin rose with nicotinic acid therapy. The daily profile of plasma glucose during hospitalization revealed an overall 16% elevation in mean plasma glucose levels during the nicotinic acid period. Finally, treatment with nicotinic acid induced marked glycosuria in some patients. Two-hourly profiles of plasma glucose on the last day of hospitalization also revealed elevated plasma glucose values during nicotinic



acid therapy. Therefore, it can be argued that the benefits of improving lipoprotein values during administration of nicotinic acid to patients with NIDDM may be counterbalanced by worsening hyperglycemia.

The results of our study suggest that a number of patients whose hyperglycemia is well controlled by dietary therapy alone may need to take hypoglycemic agents during nicotinic acid treatment. In others, the dosage of insulin or oral hypoglycemic drugs may have to be increased to control nicotinic acid-induced hyperglycemia. There are theoretical objections to increasing the insulin dosage for correction of metabolic derangements caused by another agent. For example, marked hyperinsulinemia may have a direct role in atherogenesis.<sup>22</sup> Furthermore, modest increases in insulin dosage may not be able to correct nicotinic acid-induced hyperglycemia, as was observed in one of the patients. Thus, it cannot be assumed that the worsening of hyperglycemia with nicotinic acid can be easily corrected by increasing the dosage of insulin or oral hypoglycemic drugs. Since the hyperglycemic action of nicotinic acid may be dose-dependent, some may argue that the dosage of nicotinic acid can be reduced if glycemic control deteriorates. However, the improvement in the lipoprotein profile likewise may not be optimal.

The mechanism for the hyperglycemic action of nicotinic acid in patients with NIDDM is not clear. Recently, it has been reported that nicotinic acid therapy may induce insulin resistance in normal, healthy volunteers.<sup>23</sup> The same could be true for patients with NIDDM. Whether nicotinic acid has any adverse effects on beta-cell function is not known, but there is no evidence to support such an action.<sup>15,22</sup> Another possibility is that, by interfering with triglyceride synthesis in the liver, nicotinic acid may enhance utilization of fatty acids at the expense of glucose; if so, this could lead to enhanced hepatic glucose output, another potential cause of hyperglycemia.

Another adverse effect of nicotinic acid therapy in this study was a consistent increase in plasma uric acid levels. Long-term therapy with nicotinic acid is known to increase the occurrence of acute gouty arthritis and to require greater usage of antigout medication.<sup>24</sup> Since patients with impaired glucose tolerance and NIDDM may be predisposed to develop hyperuricemia and gout,<sup>25,26</sup> nicotinic acid therapy may further increase the risk for development of gout. Although not all investigators agree that asymptomatic hyperurice-

mia is nephrotoxic, increases in plasma uric acid levels may not be benign in patients with NIDDM who are predisposed to diabetic nephropathy. Indeed, in two of our patients, marked hyperuricemia caused by nicotinic acid therapy further compromised their renal function.

To summarize, nicotinic acid therapy markedly improves the lipoprotein profile of patients with NIDDM. Although nicotinic acid generally was well tolerated in the current patients, it is known to have a variety of side effects that preclude its use in many patients. For patients with NIDDM in particular, two side effects emerge as especially worrisome. First, the drug causes deterioration of glucose control, which, for long-term therapy, must be considered a definite drawback. Also, nicotinic acid raises uric acid levels, which increases the risk for gout and could have a negative effect on renal function. For most patients with NIDDM who have dyslipidemia, therefore, nicotinic acid therapy must be used with caution, although it may be useful in primary forms of dyslipidemia. On the basis of our previous studies, we suggest that a hydroxymethylglutaryl coenzyme A reductase inhibitor<sup>27</sup> or, for marked hypertriglyceridemia, a fibric acid derivative<sup>28</sup> may be preferable as a lipid-lowering drug. Further studies, however, are needed to identify the optimal pharmacologic approach to lipid lowering in patients with NIDDM.<sup>29</sup>

This study was supported in part by the Veterans Administration, by grant HL-29252 from the National Institutes of Health, by the Southwestern Medical Foundation, and by the Moss Heart Foundation of Dallas, Tex.

We thank Raymond Wheatley, MS, Laura Caldwell, Virginia Harris, Marjorie Whelan, RN, and the nursing and dietetic services of the metabolic ward of the Veterans Administration Medical Center, Dallas, for excellent technical support, and Beverley V. Adams, MS, for statistical analysis using the CLINFO program.

#### References

1. Stern MP, Patterson JK, Haffner SM, Hazuda HP, Mitchell BD. Lack of awareness and treatment of hyperlipidemia in type II diabetes in a community survey. *JAMA*. 1989;262:860-864.
2. Barrett-Connor E, Orchard T. Diabetes and heart disease. In: National Diabetes Data Group. *Diabetes in America: Diabetes Data Group. 1984*. Washington DC, US Dept of Health and Human Services; 1985:chap 16. National Institutes of Health publication 85-1468.
3. National Cholesterol Education Program Expert Panel, National Heart Lung and Blood Institute. Report of the National Cholesterol Education Program Expert Panel on detection, evaluation, and treatment of high blood cholesterol in adults. *Arch Intern Med*. 1988;148:36-69.
4. National Diabetes Data Group. Classification and diagnosis of diabetes mellitus and other categories of glucose tolerance. *Diabetes*. 1979;28:1039-1057.
5. American Diabetes Association. Nutritional recommendations and principles for individuals with diabetes mellitus: 1986. *Diabetes Care*. 1987;10:126-132.
6. Blankenhorn DM, Nessim SA, Johnson RL, Sanmarco ME, Azen SP, Cashin-Hemphill L. Beneficial effects of combined colestipol-niacin therapy on coronary atherosclerosis and coronary venous bypass grafts. *JAMA*. 1987;257:3233-3240.
7. *Manual of Laboratory Operations: Lipid Research Clinics Program: Lipid and Lipoprotein Analysis*. 2nd ed. Washington, DC: US Dept of Health and Human Services; 1982.
8. Roeschlau P, Berni E, Gruber W. Enzymatische Bestimmung gesamtcholesterins im Serum. *Z Klin Chem Klin Biochem*. 1974;12:226.
9. Wahlefeld AW. Triglyceride determination after enzymatic hydrolysis. In: Bergmeyer HU, ed. *Methods of Enzymatic Analysis*. Orlando, Fla: Academic Press Inc; 1974;4:1831-1835.
10. Zar JH. *Biostatistical Analysis*. 2nd ed. Englewood Cliffs, NJ: Prentice-Hall International Inc; 1984:222-226.
11. Winer BJ. *Statistical Principles in Experimental Design*. 2nd ed. New York, NY: McGraw-Hill International Book Co; 1971:261-308.
12. Altshul R, Hoffer A, Stephen JD. Influence of nicotinic acid on serum cholesterol in man. *Arch Biochem Biophys*. 1955;54:558-559.
13. Gurian H, Adlersberg D. The effect of large doses of nicotinic acid on circulating lipids and carbohydrate tolerance. *Am J Med Sci*. 1959;237:12-22.
14. Gaut ZN, Pocelinko R, Solomon HM, Thomas GB. Oral glucose tolerance, plasma insulin, and uric acid excretion in man during chronic administration of nicotinic acid. *Metabolism*. 1971;20:1031-1035.
15. Miettinen TA, Taskinen M-R, Pelkonen R, Nikkila EA. Glucose tolerance and plasma insulin in man during acute and chronic administration of nicotinic acid. *Acta Med Scand*. 1969;186:247-253.
16. Parsons WB Jr. Studies of nicotinic acid use in hypercholesterolemia: changes in hepatic function, carbohydrate tolerance and uric acid metabolism. *Arch Intern Med*. 1961;107:85-99.
17. Molnar GD, Berge KG, Rosevear JW, McGuckin WF, Achon WF. The effect of nicotinic acid in diabetes mellitus. *Metabolism*. 1964;13:181-189.
18. Vega GL, Grundy SM. Gemfibrozil therapy in primary hypertriglyceridemia associated with coronary heart disease: effects on metabolism of low-density lipoproteins. *JAMA*. 1986;255:2398-2403.
19. Garg A, Grundy SM. Gemfibrozil alone and in combination with lovastatin for treatment of hypertriglyceridemia in NIDDM. *Diabetes*. 1989;38:364-372.
20. Simons LA, Hickie JB, Balasubramanian S. On the effects of dietary n-3 fatty acids (Maxepa) on plasma lipids and lipoproteins in patients with hyperlipidemia. *Atherosclerosis*. 1985;54:75-88.
21. Sullivan DR, Sanders TAB, Trayner IM, Thompson GR. Paradoxical elevation of LDL apoprotein B levels in hypertriglyceridemic patients and normal subjects ingesting fish oil. *Atherosclerosis*. 1986;61:129-134.
22. Stout RW. Insulin and atheroma: 20-year perspective. *Diabetes Care*. 1990;13:631-654.
23. Kahn SE, Beard JC, Schwartz MW, et al. Increased B-cell secretory capacity as mechanism for islet adaptation to nicotinic acid-induced insulin resistance. *Diabetes*. 1989;38:562-568.
24. Coronary Drug Project Research Group. Clofibrate and niacin in coronary heart disease. *JAMA*. 1975;231:360-381.
25. Weiss TE, Segaloff A, Moore C. Gout and diabetes. *Metabolism*. 1957;6:103-106.
26. Whitehouse FW, Cleary WJ Jr. Diabetes mellitus in patients with gout. *JAMA*. 1966;197:73-76.
27. Garg A, Grundy SM. Lovastatin for lowering cholesterol levels in non-insulin-dependent diabetes mellitus. *N Engl J Med*. 1988;318:81-86.
28. American Diabetes Association. Role of cardiovascular risk factors in prevention and treatment of macrovascular disease in diabetes. *Diabetes Care*. 1989;12:573-579.

## Lym Sevi

David A. Lea

The clinical important di severe hype lymphadenc should be in

THE CLINI  
ized lymphac  
ened by its p  
acquired im  
and the lyn  
but it is impc  
aware of othe  
the different  
triglyceriden  
has been assc  
omas, abdom  
hepatomegal  
thy, chronic  
foam-cell infil  
polyneuropat  
uricemia<sup>24</sup>; it  
scribed as a c  
pathy. We no  
be the first c  
denopathy w  
severe hyper  
micronemia (1

#### Report of a C

A 47-year-  
diabetic won  
Oregon Hes  
Hospital, P  
pain, severe l  
na triglycer  
>2000 mg/dl  
She was dis  
ses mellitus at  
y of lymph  
eridemia is n  
ague histori  
ing aware of  
nodes even w  
ood health s  
claimed to ha  
illary lymph  
years old, bu  
was unavaila  
age 35 year  
en, bilateral

From the Division of Medicine, University of California at Los Angeles (Dr. Lea), and the Division of Endocrinology and Metabolic Diseases, Division of Medicine, A-15, Veterans Ave, Los Angeles, Calif 90073.

U.S. copyright law (title 17 of U.S. code) governs the reproduction and redistribution of copyrighted material.